

Letters to the Editor

Indian J. Phys. **49**, 401-402 (1975)

On the solution of equations of connection in the new unified field theory

N. N. GHOSH

S. N. Bose Institute of Physical Sciences, 92, Acharya Prafulla Chandra Road, Calcutta 700009

(Received 5 December 1974)

In the previous note (Ghosh 1973) the equations of connection were obtained as

$$\partial_\sigma S_{\mu\nu} = S_{\lambda\nu} P^\lambda_{\mu\sigma} - F_{\lambda\nu} V^\lambda_{\mu\sigma} + S_{\lambda\mu} P^\lambda_{\nu\sigma} - F_{\lambda\mu} V^\lambda_{\nu\sigma} \quad \dots (1)$$

$$\partial_\sigma F_{\mu\nu} = -S_{\lambda\nu} V^\lambda_{\mu\sigma} + F_{\lambda\nu} P^\lambda_{\mu\sigma} + S_{\lambda\mu} V^\lambda_{\nu\sigma} - F_{\lambda\mu} P^\lambda_{\nu\sigma} \quad \dots (2)$$

where

$$S_{\mu\nu} = \frac{1}{2}(g_{\mu\nu} + g_{\nu\mu}), \quad F_{\mu\nu} = \frac{1}{2}(g_{\mu\nu} - g_{\nu\mu}),$$

$$P^\lambda_{\mu\nu} = \frac{1}{2}(\tilde{\Gamma}^\lambda_{\mu\nu} + \Gamma^\lambda_{\mu\nu}), \quad V^\lambda_{\mu\nu} = \frac{1}{2}(\tilde{\Gamma}^\lambda_{\mu\nu} - \Gamma^\lambda_{\mu\nu}).$$

The 80 connection coefficients $\Gamma^\lambda_{\mu\nu}$, $\tilde{\Gamma}^\lambda_{\mu\nu}$ symmetric in μ, ν are restricted by the conditions

$$\Gamma^\lambda_{\mu\mu} = \tilde{\Gamma}^\lambda_{\mu\mu} \quad (\mu \text{ not summed}) \quad \dots (3)$$

$$\tilde{\Gamma}^\lambda_{\mu\nu} \approx \Gamma^\lambda_{\mu\nu}.$$

From eq. (1) we obtain

$$S^\rho_{\mu\nu} = P^\rho_{\mu\nu} - S^{\rho\sigma} F_{\lambda\sigma} V^\lambda_{\mu\nu}. \quad \dots (4)$$

By substituting eq. (4) in eq. (2) we have

$$\partial_\sigma F_{\mu\nu} - F_{\rho\nu} S^\rho_{\mu\sigma} + F_{\rho\mu} S^\rho_{\nu\sigma} = (S_{\lambda\mu} - F_{\rho\mu} S^{\rho\sigma} F_{\lambda\sigma}) V^\lambda_{\nu\sigma} - (S_{\lambda\nu} - F_{\rho\nu} S^{\rho\sigma} F_{\lambda\sigma}) V^\lambda_{\mu\sigma} \quad \dots (5)$$

Denoting

$$S_{\lambda\mu} - F_{\rho\mu} S^{\rho\sigma} F_{\lambda\sigma} \text{ by } U_{\lambda\mu}$$

and noting $U_{\lambda\mu} = U_{\mu\lambda}$, eq. (5) is expressed as

$$F_{\mu\nu\sigma} = U_{\lambda\mu} V^\lambda_{\nu\sigma} - U_{\lambda\nu} V^\lambda_{\mu\sigma}, \quad \dots (6)$$

where

$$F_{\mu\nu\sigma} + F_{\nu\sigma\mu} + F_{\sigma\mu\nu} = 0. \quad \dots (7)$$

Referring to the field equations proposed in the previous note (Ghosh 1973) we have to incorporate the set of auxiliary equations further

$$V^\sigma_{\mu\sigma} = 0, \quad P^\sigma_{\mu\sigma} = \partial_\mu \log(-g)^{\frac{1}{2}}. \quad \dots (8)$$

We have now the system of eqs. (6) and (8) to determine V 's while the P 's are given by eq. (4).

Let us apply the above formulae to the case of a special non-symmetric tensor field $g_{\mu\nu}$ with symmetric components $S_{11}, S_{22}, S_{33}, S_{44}$ and the anti-symmetric components $F_{14}, F_{23}, F_{32}, F_{41}$, where all the components are functions of four variables v_1, v_2, v_3, v_4 . In the non-symmetric field theory of Einstein this case has been tackled in a more complicated way to obtain the solution of non-symmetric Γ 's (Ghosh 1955).

We first note that the surviving components of $U_{\lambda\mu}$ in eq. (6) are

$$\begin{aligned} U_{11} &= S_{11} (1 - C_4^1 C_1^4), & U_{22} &= S_{22} (1 - C_3^2 C_2^3), \\ U_{33} &= S_{33} (1 - C_3^2 C_2^3), & U_{44} &= S_{44} (1 - C_4^1 C_1^4), \end{aligned} \quad (9)$$

where $C^\lambda{}_\mu$ denotes $F_{\lambda\mu}/S_{\lambda\lambda}$.

From eq. (3) one gets $V_{\mu\mu}{}^\lambda = 0$, (μ , not summed). Using the above we now frame from eq. (6) the following set of equations

$$\begin{aligned} F_{1v1} &= U_{11} V_{v1}{}^1, & F_{3v3} &= U_{33} V_{v3}{}^3, \\ F_{2v2} &= U_{22} V_{v2}{}^2, & F_{4v4} &= U_{44} V_{v4}{}^4. \end{aligned} \quad (10)$$

Rest of the equations appear as follows

$$\begin{aligned} F_{123} &= U_{11} V_{23}{}^1 - U_{22} V_{13}{}^2, & F_{134} &= U_{11} V_{34}{}^1 - U_{33} V_{14}{}^3, \\ F_{231} &= U_{22} V_{31}{}^2 - U_{33} V_{21}{}^3, & F_{341} &= U_{33} V_{41}{}^3 - U_{44} V_{31}{}^4, \\ F_{124} &= U_{11} V_{24}{}^1 - U_{22} V_{14}{}^2, & F_{234} &= U_{22} V_{34}{}^2 - U_{33} V_{24}{}^3, \\ F_{241} &= U_{22} V_{41}{}^2 - U_{44} V_{21}{}^4, & F_{342} &= U_{33} V_{42}{}^3 - U_{44} V_{32}{}^4. \end{aligned} \quad (11)$$

To solve for P 's we have now the system of eq. (4)

$$\begin{aligned} P_{\mu\mu}{}^\lambda &= S_{\mu\mu}{}^\lambda = \Gamma_{\mu\mu}{}^\lambda, \\ P_{\mu\nu}{}^1 &= S_{\mu\nu}{}^1 - C_4^1 V_{\mu\nu}{}^4, & P_{\mu\nu}{}^3 &= S_{\mu\nu}{}^3 - C_2^3 V_{\mu\nu}{}^2, \\ P_{\mu\nu}{}^2 &= S_{\mu\nu}{}^2 - C_3^2 V_{\mu\nu}{}^3, & P_{\mu\nu}{}^4 &= S_{\mu\nu}{}^4 - C_1^4 V_{\mu\nu}{}^1. \end{aligned} \quad (12)$$

It appears that in general four of the V 's remain undetermined, so also four of the P 's corresponding to them.

REFERENCES

- Ghosh N. N. 1955 *Prof. Theor. Phys.* **13**, 587.
Ghosh N. N. 1973 *Ind. J. Phys.* **47**, 187.